

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ



الكلمة الطيبة صدقة





بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

PEDIATRICS Mcq Of The Day
August.2008

Dr.Kamel Youssef Hassan
Pediatrician Consultant
C.E.S de Pédiatrie
C.P.B in Pediatrics
MRCBCH
Gaza-Palestine
E-mail: kyh10557@yahoo.com





PEDIATRICS Meg Of The Day
August, 2008





01-08-2008

Question : 1

All of the following are characteristic features of Kwashiorkor except?

A-High blood osmolarity.

B-Hypoalbuminemia.

C-Oedema.

D-Fatty liver.

THE CORRECT ANSWER is (A)

02-08-2008

Question : 2

A3 years old boy presents with poor urinary stream, .Most likely cause is?

A-Stricture urethra.

B-Neurogenic bladder.

C-Urethral calculus.

D-Posterior urethral valves.

THE CORRECT ANSWER is (D)

03-08-2008

Question : 3

A 11 months old boy, weighing 3 kg, has polyuria, polydipsia and delayed motor milestones,

blood investigations shows:

Creatinine of 0.5 mg/dl

Potassium 3 mEq/l

Sodium 125 mEq/l

Chloride 88 mEq/l

Calcium 8,8 mg/dl

Ph 7.46 , Bicarbonate 26 mEq/l

Ultrasonography shows: Medullary Nephrocalcinosis, the most likely diagnosis is ?

A-Renal tubular acidosis

B-Diabetes insipidus.

C-Bartters syndrome.

D-Pseudohypoaldosteronism.

THE CORRECT ANSWER is (C)



04-08-2008

Question : 4

The treatment of choice for primary grade V Vesicoureteric reflux involving both kidneys in a 6 months old boy is?

A-Antibiotic prophylaxis.

B-Ureteric reimplantation.

C-Cystoscopy followed by subureteric.

D-Bilateral ureterostomies.

THE CORRECT ANSWER is (B)

05-08-2008

Question : 5

The most common leucocystoclastic vasculitis affecting children is?

A-Takayasu disease.

B-Mucocutaneous lymph node syndrome.(Kawasaki disease).

C-Henoch schonlein purpura.

D-Polyarteritis nodosa.

THE CORRECT ANSWER is (C)

06-08-2008

Question : 6

Common ocular manifestation in trisomy 13 is ?

A-Capillary hemangioma.

B-Bilateral microphthalmos.

C-Neurofibroma.

D-Dermoid cyst.

THE CORRECT ANSWER is (B)

07-08-2008

Question : 7

Blood specimen for neonatal thyroid screening is obtained on?

A-Cord blood.

B-24 hours after birth.

C-48 hours after birth.

D-72 hours after birth.

THE CORRECT ANSWER is (A)



08-08-2008

Question : 8

A child with recurrent urinary tract infection is most likely to show?

A-Posterior urethral valves.

B-Vesicoureteric reflux.

C-Neurogenic bladder.

D-Renal and Ureteric calculi.

THE CORRECT ANSWER is (B)

09-08-2008

Question : 9

The appropriate approach to a neonate presenting with vaginal bleeding on day 4 of life is?

A-Administration of vit K.

B-Investigation for bleeding disorder.

C-No specific therapy.

D-Administration of 10ml/kg of fresh frozen plasma over 4 hours.

THE CORRECT ANSWER is (C)

10-08-2008

Question : 10

Which one of the following drugs is used for foetal therapy of congenital adrenal hyperplasia?

A-Hydrocortisone.

B-Prednisolone.

C-Fludrocortisone.

D-Dexamethasone.

THE CORRECT ANSWER is (D)

11-08-2008

Question : 11

The coagulation profile in a 13 years old girl with menorrhagia having Von Willebrand disease is?

A-Isolated prolonged PTT with a normal PT.

B-Isolated prolonged PT with a normal PTT.

C-Prolongation of both PT and PTT.

D-Prolongation of thrombin time.

THE CORRECT ANSWER is (A)



12-08-2008

Question : 12

All of the following are true about manifestations of vitamin E deficiency, except?

- A-Hemolytic Anemia.
- B-Posterior column abnormalities.
- C-Cerebellar ataxia.

D-Autonomic dysfunction.

THE CORRECT ANSWER is (D)

13-08-2008

Question : 13

All of the following therapies may be required in a 1 hour old infant with sever birth asphyxia except?

- A-Glucose.
- B-Dexamethasone.**
- C-Calcium gluconate.
- D-Normal saline.

THE CORRECT ANSWER is (B)

14-08-2008

Question : 14

The most common cause of renal scaring in a 3 years old child's

- A-Trauma
- B-Tuberculosis
- C-Visicuureteral reflux indused pyelonephritis**
- D-Intersteial nephritis

THE CORRECT ANSWER is (C)

15-08-2008

Question : 15

Which one of the following is the common cause of congenital hydrocephalus

- A-Craniosynostosis
- B-Intrauterine meningitis
- C-Aqueductal stenosis**
- D-Malformation of great vein of Galen

THE CORRECT ANSWER is (C)



16-08-2008

Question : 16

In a child, non functioning kidney is best diagnosed by

A-Ultrasonography

B-IVU

C-DTPA renogram

D-Creatinine clearance

THE CORRECT ANSWER is (C)

17-08-2008

Question : 17

The most common malignant neoplasm of infancy is

A-Malignant teratoma

B-Neuroblastoma

C-Wilm's tumor

D-Hepatoblastoma

THE CORRECT ANSWER is (B)

18-08-2008

Question : 18

The most common presentation of a child with Wilm's tumor is

A-An asymptomatic abdominal mass

B-Haematuria

C-Hypertension

D-Hemoptysis due to pulmonary secondary

THE CORRECT ANSWER is (A)

19-08-2008

Question : 19

Eisenmenger's syndrome is characterised by all except

A-Return of left ventricle and right ventricle to normal size

B-Pulmonary veins not distended

C-Pruning of peripheral pulmonary arteries

D-Dilatation of central pulmonary arteries

THE CORRECT ANSWER is (A)



20-08-2008

Question : 20

The most common site of leak in CSF rhinorrhea is

A-Sphenoid sinus

B-Frontal sinus

C-Cribriform plate

D-Tegmen tympani

THE CORRECT ANSWER is (C)

21-08-2008

Question : 21

Which of the following is the most common renal cystic disease in infants is.

A-Polycystic kidney

B-Simple renal cyst

C-Unilateral renal dysplasia

D-Calyceal cyst

THE CORRECT ANSWER is (C)

22-08-2008

Question : 22

The most common type of total anomalous pulmonary venous connection is

A-Supracardiac

B-Infracardiac

C-Mixed

D-Cardiac

THE CORRECT ANSWER is (A)

23-08-2008

Question : 23

Diagnosis of beta Thalassemia is established by

A-NESTROFT test

B-HbA1 C estimation

C-Hb electrophoresis

D-Target cells in peripheral smear

THE CORRECT ANSWER is (C)



24-08-2008

Question : 24

A child having status epilepticus should not be given

A-Clonazepam

B-Diazepam

C-Lamotrigine

D-Phosphenytoin

THE CORRECT ANSWER is (C)

25-08-2008

Question : 25

The sodium content of ReSoMal (rehydrarion solution for malnourished children is ?

A-90 mmol/L

B-60 mmol/L

C-45mmol/L

D-30 mmol/L

THE CORRECT ANSWER is (C)

26-08-2008

Question : 26

Which of the following statements is true of primary grade IV-V Vesicoureteric reflux in young children?

A-Renal scaring usually begins in the midpolar regions

B-Postnatal scaring may occur even in the absence of urinary tract infections

C-Long term outcome is comparable in patients treated with either antibiotic prophylaxis or surgery.

D-Oral amoxicillin is the choice antibiotic for prophylaxis

THE CORRECT ANSWER is (B)



27-08-2008

Question : 27

A 3 years old boy presents with fever, dysuria and gross hematuria. Physical examination shows a prominent suprapubic area which is dull to percussion. Urinalysis reveals red blood cells but no Proteinuria.

Which of the following is the most likely diagnosis?

A-Acute Glomerulonephritis

B-Urinary tract infection

C-posterior urethral valves

D-Teratoma.

THE CORRECT ANSWER is (B)

28-08-2008

Question : 28

Transient synovitis (Toxic synovitis) of the hip is characterized by all of the following except ?

A-May follow upper respiratory infection.

B- ESR and W.B.C counts are usually normal.

C-ultrasound of the joint reveals widening of the joint space.

D-The hip is typically held in adduction and internal rotation.

THE CORRECT ANSWER is (D)

29-08-2008

Question : 29

Enzyme replacement therapy is available for which of the following ?

A-Gaucher disease

B-Niemann Pick disease.

C-Mucopolidosis.

D-Metachromatic leukodystrophy

THE CORRECT ANSWER is (A)



30-08-2008

Question : 30

Cardiomyopathy may be seen in all of the following except

- A-Duchene muscular dystrophy.
- B-Friedreich's ataxia.
- C-Type II glycogen storage disease.

D-Alkaptonuria

THE CORRECT ANSWER is (D)

31-08-2008

Question : 31

All of the following are true of beta thalassemia major except?

- A-Splenomegaly
- B-Target cells on peripheral smear.
- C-Microcytic hypochromic anaemia.

D-Increased osmotic fragility.

THE CORRECT ANSWER is (D)



